arXiv:1412.2614v1 [math-ph] 8 Dec 2014

On eigenfunctions of one-dimensional Schrödinger operator with polynomial potentials *

Andrey E. Mironov †, Bayan T. Saparbaeva ‡

One-dimensional Schrödinger operator with polynomial potentials of degree 3 and 4 appears in many areas of mathematical physics (see e.g. [1]–[3]). In this paper we point out an connection between eigenfunctions of such operator and eigenfunctions of commuting ordinary differential operators of rank two (rank two Baker–Akhiezer function).

We recall that if differential operators $L_n = \sum_{i=0}^{n} u_i(x)\partial_x^i$ and $L_m = \sum_{j=0}^{m} v_j(x)\partial_x^j$ commute, then by Burchnall–Chaundy lemma [4] there exist a non-zero polynomial $R(z, w)$ such that

*The work was supported by RSF (grant 14-11-00441).

†Sobolev Institute of Mathematics, Ac. Koptyug avenue 4, 630090, Novosibirsk, Russia, and Department of Mathematics and Mechanics, Novosibirsk State University, Pirogov street 2, 630090 Novosibirsk, Russia; e-mail: mironov@math.nsc.ru

‡Department of Mathematics and Mechanics, Novosibirsk State University, Pirogov street 2, 630090 Novosibirsk, Russia; e-mail: saparbayevabt@gmail.com

$R(L_n, L_m) = 0$. The polynomial $R(z, w)$ defines the *spectral curve* $\Gamma = \{(z, w) \in \mathbb{C} : R(z, w) = 0\}$. If ψ is an eigenfunction $L_n\psi = z\psi$, $L_m\psi = w\psi$, then $P = (z, w) \in \Gamma$. The *rank* of the pair L_n, L_m is $l = \dim\{\psi : L_n\psi = z\psi, L_m\psi = w\psi\}$, where (z, w) is a general point in Γ. In the case of operators of rank 1 eigenfunctions are expressed via theta-function of the Jacobi variety of Γ (see [5]). At $l > 1$ eigenfunctions correspond to the *spectral data* (see [6]) $S = \{\Gamma, q, k^{-1}, \gamma_1, ..., \gamma_{lg}, \beta_1, ..., \beta_{lg}\}$, where Γ is a Riemann surface of genus g, $q \in \Gamma$ is a marked point with a local parameter k^{-1}, $\gamma_1, ..., \gamma_{lg} \in \Gamma$ is a set of points and $\beta_j = (\beta_{j,1}, ..., \beta_{j,l-1})$ is a set of vectors. *Baker–Akhiezer function of rank l* $\psi(x, P) = (\psi_1, ..., \psi_l), P \in \Gamma$ is a function which satisfies the conditions:

1. On $\Gamma \setminus \{q\}$ function ψ has poles in γ_i and

$$\operatorname{res}_{\gamma_i} \psi_j(x, P) = \beta_{i,j} \operatorname{res}_{\gamma_i} \psi_l(x, P).$$

2. In the neighbourhood of q the function $\psi(x, P)$ has the form

$$\psi(x, P) = \left(\sum_{i=0}^{\infty} \xi_i(x) k_j^{-i} \right) \Psi(x, k),$$

where $\Psi(x,P)$ is a solution of the equation $\Psi_x = A\Psi$,

$$A = \begin{pmatrix} 0 & 1 & 0 & ... & 0 & 0 \\ 0 & 0 & 1 & ... & 0 & 0 \\ ... & ... & ... & ... & ... & ... \\ 0 & 0 & 0 & ... & 0 & 1 \\ k+\omega_1(x) & \omega_2(x) & \omega_3(x) & ... & \omega_{l-1}(x) & 0 \end{pmatrix},$$

$\omega_j(x)$ are functional parameters. For the spectral data in general position there is a unique Baker–Akhiezer function. Herewith for a meromorphic function $f(P)$ on Γ with the pole in q of order n there is a unique operator L_f of order ln such that $L_f\psi = f(P)\psi$. For such meromorphic functions f and g operators L_f and L_g commute. At $l > 1$ Baker–Akhiezer function can not be found explicitly. Nevertheless the operators themselves can be found in some cases. The operators of ranks two and three in the case of elliptic spectral curves were found in [6], [7]. A new method of finding rank two operators corresponding to hyperelliptic spectral curves were suggested in [8] (see also [9]–[12]). In particular, in [8] it is proved that $L_4 = (\partial_x^2 + \alpha_3 x^3 + \alpha_2 x^2 + \alpha_1 x + \alpha_0)^2 + \alpha_3 g(g+1)x$ commutes with L_{4g+2}. Operators L_4, L_{4g+2} define commutative subalgebra in the first Weyl algebra. At $g = 1$ these operators firstly appeared in [13]. It is turn out that there is a relation between eigenfunctions of L_4 at $g = 2, 4$ and functions from the kernel

of $L_2 = \partial_x^2 + \alpha_3 x^3 + \alpha_2 x^2 + \alpha_1 x + \alpha_0$. Let φ be a solution of $L_2\varphi = 0$.

Theorem 1 *1. Let $g = 2$, z be a solution of the equation*

$$z^2 + 4\alpha_2 z + 12\alpha_1\alpha_3 = 0.$$

Then $L_4\psi = z\psi$, where $\psi = p\varphi$, $p(x) = 6\alpha_3 x + z + 4\alpha_2$.
2. Let $g = 4$, z be a solution of the equation

$$z^3 + 20\alpha_2 z^2 + 16(4\alpha_2^2 + 13\alpha_1\alpha_3)z + 320\alpha_3(7\alpha_0\alpha_3 + 2\alpha_1\alpha_2) = 0.$$

Then $L_4\psi = z\psi$, where $\psi = p\varphi$, $p(x) = 280\alpha_3^2 x^2 + 20\alpha_3(z + 16\alpha_2)x + z^2 + 20\alpha_2 z + 64\alpha_2^2 + 168\alpha_1\alpha_3$.

Let $l_4 = p^{-1}L_4 p$, $l_{4g+2} = p^{-1}L_{4g+2}p$ (the function $p(x)$ is pointed out in Theorem 1).

Corollary 1 *At $g = 2, 4$ operators l_4, l_{4g+2}, L_2 form a commutative ring modulo L_2, i.e.*

$$[l_4, L_2] = B_1 L_2, \quad [l_{4g+2}, L_2] = B_2 L_2,$$

where B_1, B_2 are some operators.

Using [8] one can show that for $\alpha_3^3 - 4\alpha_2\alpha_3\alpha_4 + 8\alpha_1\alpha_4^2 = 0$ the operator

$$L_4^a = (\partial_x^2 + \alpha_4 x^4 + \alpha_3 x^3 + \alpha_2 x^2 + \alpha_1 x + \alpha_0)^2 + 2g(g+1)x(\alpha_3 + 2\alpha_4 x)$$

commutes with an operator of order $4g+2$ (in a partial case it was observed by V. Oganesyan [14]). Let φ be a solution of the equation $(\partial_x^2+\alpha_4x^4+\alpha_3x^3+\alpha_2x^2+\alpha_1x+\alpha_0)\varphi=0$.

Theorem 2 *1. Let $g=1$, $z=\frac{\alpha_3^2}{\alpha_4}-4\alpha_2$. Then $L_4^a\psi=z\psi$, where $\psi=p\varphi,\ p(x)=4\alpha_4x+\alpha_3$.*
2. Let $g=2$, z be a solution of the equation

$$z^2-\left(\frac{3\alpha_3^2}{\alpha_4}-16\alpha_2\right)z+24\alpha_1\alpha_3+192\alpha_0\alpha_4=0.$$

Then $L_4^a\psi=z\psi$, where $\psi=p\varphi$, $p(x)=24\alpha_4^2x^2+12\alpha_3\alpha_4x-3\alpha_3^2+\alpha_4(z+16\alpha_2)$.

The operator $L_4^b=(\partial_x^2+\alpha_1e^x+\alpha_0)^2+\alpha_1g(g+1)e^x$ commutes with an operator of order $4g+2$ [12]. Let us denote by φ a solution of the equation $\left(\partial_x^2+\alpha_1e^x+\alpha_0+\frac{1}{4}(g+\varepsilon)^2\right)\varphi=0$.

Theorem 3 *1. Let $\varepsilon=0$, then $L_4^b\psi=-\frac{1}{4}g^2(4\alpha_0+g^2)\psi$, where $\psi=p\varphi$, $p(x)=e^{gx/2}$.*
2. Let $\varepsilon=1$, then $L_4^b\psi=-\frac{1}{4}(g+1)^2(4\alpha_0+(g+1)^2)\psi$, where $\psi=p\varphi$, $p(x)=e^{-(g+1)x/2}$.

Remark 1 *Solutions of the equation $(\partial_x^2+\alpha_1e^x+\alpha_0)\varphi=0$ are expressed via Bessel function, namely the change of the variable $x=\ln\left(\frac{y^2}{4\alpha_1}\right)$ reduces this equation to $(y^2\partial_y^2+y\partial_y+(y^2+4\alpha_0))\varphi=0$.*

The authors are grateful to B.A. Dubrovin for valuable discussions.

References

[1] *Masoero D.* // J. Phys. A: Math. Theor. 2010. V. 43. P. 1–28.

[2] *Chudnovsky D., Chudnovsky G.* // Acta Applicandae Mathematicae 1994. V. 279. P. 167–185.

[3] *Dorey P., Tateo R.* // Nucl. Phys. 1999. V. B563. P. 573–602.

[4] *Burchnall J.L., Chaundy T.W.* // Proc. London Math. Soc. Ser. 2 1923. V.21. P. 420–440.

[5] *Krichever I.M.* // Functional Anal. Appl. 1977. V. 11. N. 1. P. 12–26.

[6] *Krichever I.M., Novikov S.P.* // Russian Math. Surveys 1980. V.35. N. 6. P. 47–68.

[7] *Mokhov O.I.* // Mathematics of the USSR-Izvestiya. 1990. V. 35. N. 3. P. 629–655.

[8] *Mironov A.E.* // Invent math 2014. V.197. N.2. P. 417–431.

[9] *Mironov A.E.* // Amer. Math. Soc. Transl. Ser. 2. 2014. V. 234. 309–322.

[10] *Mironov A.E.* // Proceedings of 6th European Congress of Mathematics. 2013. P. 459–473.

[11] *Mironov A.E.* // Funct. Anal. Appl. 2005. V. 39. N. 3. P. 240–243.

[12] *Dabletshina V.N.* // Siberian Electronic Math. Reports. 2013. V. 10. P. 109–112.

[13] *Dixmier J.* // Bull. Soc. Math. France 1968. V.96. P.209–242.

[14] *Oganesyan V.* // arXiv:1409.4058.